

Review Article

High Intensity Interval Training or Moderate Intensity Continuous Exercise in Patients with Myocardial Infarction?

Irene C. Themistocleous and Manos Stefanakis

Department of Life & Health Sciences, University of Nicosia, 46, Makedonitissas Avenue, Nicosia, 1700, Cyprus

Abstract

Objectives: To determine the effects of high intensity interval training (HIIT) or moderate intensity continuous exercise (MICE) in patients with myocardial infarction (MI). **Background:** Cardiovascular diseases are the leading cause of mortality and morbidity globally causing a significant reduction in the quality of life of these patients. Participation of these patients in rehabilitation programs which involve a significant component of exercise seems to help by improving functional capacity and quality of life (QoL). Despite the beneficial effect of exercise, the type of exercise that yields the best results is yet to be determined. **Methods:** Three databases (MEDLINE, CINHALL and SportDirect) were searched in May-June 2017 for original articles regarding the effect of two types of exercise in patients with myocardial infarction. Randomized control trial studies which enrolled patients with myocardial infarction and studied the effects of HIIT and/or MICE, were included in this review. Data were extracted and summarised and all studies were assessed for bias. **Results:** Both forms of exercise seem to improve the relevant outcome measures such as functional capacity, QoL, walking distance, fatigue and function of the left heart. However HIIT seemed to be better in comparison with MICE in improving these outcomes. Several limitations and risk of bias have been identified and reported. **Conclusion:** Both HIIT and MICE are recommended for patients with MI. Further research is required to conclusively support the superiority of HIIT over other types of exercise.

Corresponding Author:
Manos Stefanakis; email:
manos.stefanakis.gr@gmail.com

Production and Hosting by
Knowledge E

© Irene C. Themistocleous and Manos Stefanakis. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editor-in-Chief:
Dr. Dimitrios Papandreou
Official Publication of Zayed University, UAE



 OPEN ACCESS

1. Introduction

According to World Health Organisation [1] chronic diseases are among the primary causes of global mortality. The most common causes of mortality in 2012 were: cardiovascular diseases (46.2% of all deaths), cancer (21.7%), respiratory diseases (10.7%) and diabetes (4%) [1].

Several studies show that frequent physical exercise helps in the prevention of several chronic diseases and reduces the mortality rate from all causes, including cardiovascular diseases [2–7]. Aerobic exercise is extremely important for patients with CHD with or without MI [8]. Taylor and colleagues [9] report that rehabilitation programs based on aerobic exercise have reduced the total mortality rate by 20% and the cardiovascular disease mortality rate by 26% and they have helped in reducing risk factors such as hypertension, hypercholesterolemia, diabetes, and obesity in patients with MI, angina and CHD [9].

In addition exercise improves well-being and QoL [11]. Exercise is important especially for patients with coronary heart disease (CHD) [10] because it improves functional capacity, reduces coronary ischemia and angina and improves endothelial function [13]. However benefits from exercise rehabilitation programs depend heavily on the parameter of the program most notably the intensity and volume of exercise [12]. Studies show that perhaps interval exercise especially with high intensity might be superior in comparison with other forms of exercise in improving cardiac function, functional capacity and ultimately QoL in patients with CHD especially after myocardial infarction [14, 16], (Nillson Westheim & Risberg, 2008b; Moholdt Madssen Rognum & Aamot 2014).

2. Methods

2.1. Search Strategy

The following electronic databases were searched from March until June 2014: Science Direct, MEDLINE and Cumulative Index to Nursing and Allied Health Literature [CINAHL]. The medical subject headings (MeSH) “myocardial infarction”, “coronary heart disease”, “high intensity interval training” AND “moderate intensity continuous exercise” were used alone or in combination. Each specific phrase was combined with the Boolean operator to limit the search and make it more specific. Additional searches were carried out by scanning the reference lists of related articles in order to maximize the amount of investigations involved in the current review.

3. Study Criteria and Selection

In order to define and frame the research questions the Population(s), Intervention(s), Comparator(s), Outcome(s) and Study Design (PICOS) were used [17]. The titles and abstracts of all records were screened initially against the basic initial eligibility criteria. A single failed eligibility criterion was sufficient for a study to be excluded from a

Inclusion Criteria	Exclusion Criteria	PICOS
Was the study a RCT?	Study designs other than RCT Quasi experimental studies, pilot studies	Study
Was the study published in English?	Duplicates Abstract and pilot-studies publications	
Did the study include males and females with CHD and MI?	Included subjects with disease risk factors	Population
Was the intervention HIIT or MICE and was it combined with another type of exercise?	Not HIIT or MICE Not specified	Intervention
Were the outcomes measures relevant?	Not relevant outcomes measures	Outcomes
Were the interventions compared with another type of exercise, or placebo or non therapy?		Comparison

TABLE 1: Full eligibility screening criteria.

review [17]. Moreover, the records that remained after initial eligibility screening, were therefore screened against the full eligibility criteria outlined in Table 1.

3.1. Outcome Measures

All relevant outcome measures were considered. Specifically measures such as QoL, VO₂peak, functional capacity, fatigue or perceived rate of exertion, left ventricular function, and exercise time were considered relevant.

3.2. Data Extraction and Assessment of Risk of Bias

For each RCT included, data were extracted regarding: first author, year of publication, study outcomes, groups and exercise parameters, and assessed for bias using the Cochrane risk of bias tool. The quality assessment was performed to identify the quality of papers. Trials should provide full details of the randomisation process, allocation concealment or blinding of outcome assessment to be considered high quality. The quality of included studies was assessed using the Jadad scale [20], a three-point questionnaire form. Each question was to be answered with a *yes* and score a single point, or a *no* and scored zero points.

4. Results




Figure 1: PRISMA diagram for included studies [19].

4.1. Identification of Records and Study Selection

The search strategy from the three databases identified a total of 1095 records. An initial screen of the article title resulted in exclusion of 830 papers. Three hundred and sixty five studies were assessed for eligibility using the inclusion and exclusion criteria of Table 1. Following this process of the eligibility criteria, 352 records were excluded, due to reason described in detail in Figure 1. The remaining 13 records met the eligibility criteria and were included in the final review.

4.2. Description of Included Studies

Thirteen studies were included in this review. A summary of the studies' characteristics is presented in Table 2. Eight of the studies [22–26, 29–31] compared HIIT with MICE alone or in combination with a control group, whereas the remaining studies [14, 15, 21, 27, 28] examined only HIIT or MICE in relation to another type of exercise.

Six of the studies [22, 23, 25, 29–31] had a small sample (below 30) and the rest had sample that varied from 59 to 200 participants. Most of the studies used an exercise program of 10–16 weeks with a frequency of 2–3 times per week, however one study used a program of four week with a higher frequency (5times/week) and one study used a 1-year program of 2 sessions per week.

Generally both forms of exercise seemed to improve QoL, maximum aerobic capacity, functional capacity and left ventricular function. HIIT and MICE were superior to control intervention in two of the studies that had a control group [24, 31] in terms of aerobic capacity, functional capacity and QoL. In addition HIIT was superior to a control group in another two studies [14, 15] in terms of functional capacity, QoL and exercise time. HIIT was also superior in improving VO_2 peak compared to aerobic exercise (no further information regarding parameters) in two additional studies [27, 28]. No study compared MICE with a control group only. From the eight studies that examined both HIIT and MICE, five favoured HIIT [25, 26, 29–31] in at least one outcome measure or one time point, while 3 studies [22–24] showed no differences between the two forms of exercise.

4.3. Risk of Bias Assessment

The Jadad score ranged between 0 and 3 out of 5 (Table 3). One study scored 0 [25]; two studies scored 1 [24, 30], four studies scored 2 [21–23, 31] and six studies scored 3 [14, 15, 26–29]. These scores indicated that approximately half of the studies were low quality. This is due to the fact that the double-blinding criterion is not feasible in these kind of RCT within exercise intervention. On the Jadad scale 40% of the score accounts for the double-blinding criterion (Hempel et al., 2011).

All the included studies were evaluated in terms of its risk of bias (Table 4). Major sources of the risk of bias were related to allocation concealment, blinding study subjects or research personnel and blinding of outcome assessment. Risk of reporting bias was low in general. Therefore, a high risk of bias might be introduced in most of the RCTs included.

Authors	Sample size/Groups	Duration / Frequency	Equipment	Exercise protocol	Outcome measures
[29]	21 participants IT/MICE	10 weeks 3days/week	Treadmill	IT group 4x4min at 80-90%VO2peak [33min] MICE group 50-60%VO2peak [41min]	IT group had greater ↑ of VO2peak compared to MICE group
[30]	14 participants IT/MICE	16 weeks 2days/week Additional 3days/week	Treadmill, Stair-climber, Combined arm and leg cycle ergometer	IT group 40%VO2reserve (2 min) with intervals at 90%VO2reserve (2 min) [30min] 65% VO2reserve [Resistance exercise, 30min] MICE group 65% VO2reserve [30min] 65% VO2reserve [Resistance exercise, 30min]	IT and Mice groups had similar ↑of VO2 Peak IT group had greater ↑ of AT
[31]	27 participants IT/MICE/CONTROL	12 weeks 3days/week	Treadmill	IT group 50-60% HRpeak (3 min) with intervals at 90-95% HRpeak (4 min) [38min]	IT group had greater ↑ of LV remodeling, VO2peak and endothelium function compared to MICE and CONTROL groups. QOL improved both after IT and MICE exercise. No changes occurred in the CONTROL group regarding the QOL.

TABLE 2: Summary of the studies included in the review. **Abbreviations:** OUES: oxygen uptake efficiency slope; HRpeak: heart rate peak; IT: interval training; MICE: moderate intensity continuous training; LV Function: left ventricle function; QOL: quality of life; HR: heart rate; VO2: oxygen consumption; AT: anaerobic threshold; O₂P slope: oxygen pulse; ↔: Similar improvement; ↑: Greater improvement; ↓: Reduction. FMD: Flow mediated dilation.

Authors	Sample size/ Groups	Duration/ Frequency	Equipment	Exercise protocol	Outcome measures
				MICE group 70% HRpeak [47min]	
				CONTROL group (Walking and counseling) 70% HRpeak [47min]	
[14]	80 participants IT/CONTROL	16 weeks 2days/week	Aerobic dance movements (with music) and involved the use of both upper and lower extremities, including endurance, strength, and stretching exercises	IT group 3x5-10 min at RPE = 15-18 [50min]	Greater ↑ of functional capacity, workload (watts), exercise time (seconds) and QOL significantly improved for the exercise group.
				CONTROL group Without exercise	
[15]	80 participants IT/CONTROL	1year 2days/week	Aerobic dance movements (with music) and involved the use of both upper and lower extremities, including endurance, strength, and stretching exercises	IT group 3x5-10 min at RPE = 15-18 [50min]	Greater ↑ of functional capacity, workload (watts), exercise time (seconds) and QOL significantly improved for the exercise group.
				CONTROL group Without exercise	

TABLE 2: Table continued.

Authors	Sample size/ Groups	Duration/ Frequency	Equipment	Exercise protocol	Outcome measures
[26]	59 participants IT/MICE	4weeks 5days/week	Treadmill	IT group 4x4min intervals at 90% HRmax with active pauses of 3min of walking at 70%HRmax MICE group 70%HRmax [46min]	↑ Of VO ₂ peak in both groups after 4 weeks. After 6 months IT group significantly improved VO ₂ peak compared to MICE group. QOL significantly ↑ at 4weeks and 6months, with no significant difference between the groups. No changes occurred at 4 weeks regarding the blood markers and LV function. However, there was a change at 6 months on both groups.
[25]	20 participants IT/MICE	2 sessions 2 weeks difference	Cycle ergometer	IT group 2sets of 10min composed of repeated phases of 15s at 100% PPO, interspersed by 15s of passive recovery MICE group 70% PPO AEROBIC group N/A	IT protocol resulted in lower mean ventilation for a small difference in metabolic demand. Participants preferred the IT mainly because the perceived exertion measured by the Borg scale was lower.

TABLE 2: Table continued.

Authors	Sample size/ Groups	Duration/ Frequency	Equipment	Exercise protocol	Outcome measures
[27]	107 participants IT/AEROBIC	12 weeks 2days/week	Treadmill Aerobic exercises	IT group 4x4min inter-vals at 85-95% HRmax with active pauses of 3min of walking at 70% HRmax [38min] AEROBIC group N/A	IT group had greater ↑ of VO2peak. ↔ Improvement of endothelial function, QOL and blood markers between the groups.
[28]	107 participants IT/AEROBIC	12 weeks 2days/week Measurement at 6 and 30 months	Treadmill Aerobic exercises	IT group 4x4min inter-vals at 85-95% HRmax with active pauses of 3min of walking at 70% HRmax [38min] AEROBIC group N/A	After 6 and 30 months, VO2peak in both groups declined. At 30 months the improvement of VO2peak in IT group, was still significant.
[14]	22participants IT/MICE	12 weeks 3days/week	Cycle ergometer	IT group 10minX 1-min cycling intervals at 89% PPO separated by 1-min intervals at 10% PPO MICE group 58% of PPO [50min]	↔ Improvement of endothelial function, between the groups. No change in HR recovery or variability following 12 weeks of exercise

TABLE 2: Table continued.

Authors	Sample size/ Groups	Duration/ Frequency	Equipment	Exercise protocol	Outcome measures
[15]	14 participants IT/MICE	12 weeks 2days/week	Cycle ergometer	IT group 10minX 1-min cycling intervals at 88% PPO separated by 1-min intervals at 10%PPO MICE group 60% of PPO [50min]	. FMD was increased post-training with no differences between groups. There was a significant improvement in cardiorespiratory fitness following both training programs, with no group differences.
[24]	71 participants IT/MICE/CONTROL	16 weeks 3days/week	Treadmill	IT group Alternation of 90% HRpeak and 60% HRpeak every 2min [30min] MICE group 70-75% HRmax [30min] CONTROL group Without exercise	IT group had greater ↑ of O ₂ P slope. No changes of VE/VCO ₂ slope and OUES occurred between groups.

TABLE 2: Table continued.

Authors	Sample size/ Groups	Duration/Frequency	Equipment	Exercise protocol	Outcome measures
[24]	200 participants IT/CONTINUOUS	12 weeks 3days/week	Cycle ergometer	IT group 4x4min at 90-95% HRpeak with active pauses of 3min of 50-70% HRpeak CONTINUOUS group 70-75% HRpeak	↔ Improvement of endothelial function, cardiovascular risk factors and QOL.

TABLE 2: Table continued.

Potential Score					
JADAD SCORING CRITERIA	[29]	[30]	[31]	[14]	
The study described as randomized?	1	0	1	1	
Was the method of randomization described and appropriate to conceal allocation?	1	1	1	1	
Was there a description of withdrawals and drop outs?	1	0	0	1	
Was the study described Double-blinded?	0	0	0	0	
Was the method of blinding inappropriate?	0	0	0	0	
FINAL SCORE (0 – 5)	3/5	1/5	2/5	3/5	
The study described as randomized?	1	1	1	1	
Was the method of randomization described and appropriate to conceal allocation?	1	1	1	1	
Was there a description of withdrawals and drop outs?	1	1	1	1	
Was the study described Double-blinded?	0	0	0	0	
Was the method of blinding inappropriate?	0	0	0	0	
FINAL SCORE (0 – 5)	3/5	3/5	3/5	3/5	
JADAD SCORING CRITERIA	[25]	[22]	[23]	[24]	[21]
The study described as randomized?	0	0	0	1	0
Was the method of randomization described and appropriate to conceal allocation?	0	1	1	0	1
Was there a description of withdrawals and drop outs?	0	1	1	0	1
Was the study described Double-blinded?	0	0	0	0	0
Was the method of blinding inappropriate?	0	0	0	0	0
FINAL SCORE (0 – 5)	0/5	2/5	2/5	1/5	2/5

TABLE 3: Assessment of the methodological quality of the studies using the Jadad score.

5. Limitations

Jadad et al. [20] reported that RCTs of high quality, must score between 3 and 5 points on the above scale. The quality assessment of the papers demonstrated that only six studies reached the minimum score to be considered of adequate quality. However,

	1	2	3	4	5	6	7
[29]	-	-	+	+	-	+	+
[30]	+	?	-	-	+	+	+
[31]	+	+	+	-	+	+	+
[14]	+	+	+	-	+	+	+
[15]	+	+	+	-	+	+	+
[26]	-	?	+	-	+	+	+
[27]	+	-	+	-	+	+	+
[28]	+	-	-	-	+	+	+
[25]	-	-	-	-	+	+	+
[22]	-	-	-	-	-	+	+
[23]	+	-	-	-	+	+	+
[24]	-	-	-	-	+	+	+
[21]	-	+	+	+	+	+	+

TABLE 4: Risk of bias assessment. 1, Random sequence generation (selection bias); 2, Allocation concealment (selection bias); 3 Blinding of participants and personnel (performance bias); 4, Blinding of outcome assessment (detection bias); 5, Incomplete outcome data (attrition bias); 6, Selective reporting (reporting bias); 7, Other bias. + low risk of bias - high risk of bias ? unclear risk of bias.

none of the records were excluded based on the score. Therefore, a high risk of bias exists in most studies.

The comprehensive search strategy was performed in order to minimise publication bias. Nevertheless, it may not have been sufficient to prevent bias, as identifying and improving potential biases is not easy [32]. The exclusion of non-English language records may have introduced language bias, and this is another limitation of this review. The search of the records was updated during the process of the review, to prevent any time-lag biases; however it is impossible to exclude such biases. In addition, the review included only thirteen studies and most of them had a small sample. This makes the interpretation and generalisation of the results difficult.

6. Conclusion

Despite the limitations in the studies, HIIT and MICE both improve QoL and functional capacity of patients with MI and therefore are recommended. However there seems to be a slight advantage of HIIT over MICE but this is not a universal finding. Obviously there is significant heterogeneity in methodologies among the different trials. Despite this, the findings of this review suggest that HIIT is safe and may improve QoL in patients with MI. However, additional studies with higher sample sizes and improved

methodological quality are required to confirm these findings. In light of the limitations that have been reported, the results must be interpreted with caution.

References

- [1] A. Alwan, (2014). Global status report on noncommunicable diseases 2010. World Health Organization.
- [2] A. E. Bauman, "Updating the evidence that physical activity is good for health: an epidemiological review 2000–2003," *Journal of Science and Medicine in Sport*, vol. 7, no. 1, pp. 6–19, 2004.
- [3] S. N. Blair, H. W. Kohl III, C. E. Barlow, R. S. Paffenbarger Jr., L. W. Gibbons, and C. A. Macera, "Changes in physical fitness and all-cause mortality: a prospective study of healthy and unhealthy men," *Journal of the American Medical Association*, vol. 273, no. 14, pp. 1093–1098, 1995.
- [4] M. Lee, C. C. Hsieh, and R. S. Paffenbarger, "Exercise Intensity and Longevity in Men: The Harvard Alumni Health Study," *Journal of the American Medical Association*, vol. 273, no. 15, pp. 1179–1184, 1995.
- [5] M. Hulens, G. Vansant, A. L. Claessens, R. Lysens, E. Muls, and R. Rzewnicki, "Health-related quality of life in physically active and sedentary obese women," *American Journal of Human Biology*, vol. 14, no. 6, pp. 777–785, 2002.
- [6] C. E. I. Lebrun, Y. T. van der Schouw, F. H. de Jong, H. A. P. Pols, D. E. Grobbee, and S. W. J. Lamberts, "Relations between body composition, functional and hormonal parameters and quality of life in healthy postmenopausal women," *Maturitas*, vol. 55, no. 1, pp. 82–92, 2006.
- [7] M. L. Pollock, G. A. Gaesser, J. D. Butcher et al., "ACSM position stand: the recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness, and flexibility in healthy adults," *Medicine Science in Sports Exercise*, vol. 30, no. 6, pp. 975–991, 1998.
- [8] C. A. Barboza, L. Y. Rocha, C. T. Mostarda et al., "Ventricular and autonomic benefits of exercise training persist after detraining in infarcted rats," *European Journal of Applied Physiology*, vol. 113, no. 5, pp. 1137–1146, 2013.
- [9] R. S. Taylor, A. Brown, S. Ebrahim et al., "Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials," *American Journal of Medicine*, vol. 116, no. 10, pp. 682–692, 2004.
- [10] J. Perk, G. De Backer, H. Gohlke, I. Graham, and Ž. Reiner, "European Guidelines on cardiovascular disease prevention in clinical practice (version 2012) The Fifth

- Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of nine societies and by invited experts) Developed with the special contribution of the European Association for Cardiovascular Prevention Rehabilitation (EACPR," *European Heart Journal*, p. s092, 2012.
- [11] J. Kruk, "Physical activity in the prevention of the most frequent chronic diseases: An analysis of the recent evidence," *Asian Pacific Journal of Cancer Prevention*, vol. 8, no. 3, pp. 325-338, 2007.
- [12] L. Azevedo and M. Santos, "High-Intensity Intermittent Exercise Training for Cardiovascular Disease," *Journal of Novel Physiotherapies*, vol. 4, p. 1, 2014.
- [13] P. D. Thompson, "Exercise prescription and proscriptioin for patients with coronary artery disease," *Circulation*, vol. 112, no. 15, pp. 2354-2363, 2005.
- [14] B. B. Nilsson, A. Westheim, and M. A. Risberg, "Long-Term Effects of a Group-Based High-Intensity Aerobic Interval-Training Program in Patients With Chronic Heart Failure," *American Journal of Cardiology*, vol. 102, no. 9, pp. 1220-1224, 2008.
- [15] B. B. Nilsson, A. Westheim, and M. A. Risberg, "Effects of Group-Based High-Intensity Aerobic Interval Training in Patients With Chronic Heart Failure," *American Journal of Cardiology*, vol. 102, no. 10, pp. 1361-1365, 2008.
- [16] K. Meyer, C. Foster, N. Georgakopoulos et al., "Comparison of left ventricular function during interval versus steady- state exercise training in patients with chronic congestive heart failure," *American Journal of Cardiology*, vol. 82, no. 11, pp. 1382-1387, 1998.
- [17] D. O'Connor, S. Green, and J. P. T. Higgins, "Defining the review question and developing criteria for including studies. In," in *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0. [Updated March 2011]: The Cochrane Collaboration, 2011, Chapter 5, Retrieved from: www.cochrane handbook.org*, J. P. T. Higgins and S. Green, Eds., 2011.
- [18] J. P. T. Higgins and J. J. Deeks, "Chapter 7: Selecting studies and collecting data," in *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0. [Updated March 2011]: The Cochrane Collaboration, 2011*, J. P. T. Higgins and S. Green, Eds., 2011, Retrieved from <http://www.cochrane-handbook.org>.
- [19] C. Cooper, A. Selwood, M. Blanchard, Z. Walker, G. Blizard, and G. Livingston, "Abuse of people with dementia by family carers: representative cross sectional survey," *British Medical Journal*, vol. 339, no. 7694, pp. 155-157, 2009.
- [20] A. R. Jadad, R. A. Moore, D. Carroll et al., "Assessing the quality of reports of randomized clinical trials: Is blinding necessary?" *Controlled Clinical Trials*, vol. 17, no. 1, pp. 1-12, 1996.

- [21] V. M. Conraads, N. Pattyn, C. De Maeyer et al., "Aerobic interval training and continuous training equally improve aerobic exercise capacity in patients with coronary artery disease: the SAINTEX-CAD study," *International Journal of Cardiology*, vol. 179, pp. 203–210, 2015.
- [22] K. D. Currie, J. B. Dubberley, R. S. McKelvie, and M. J. MacDonald, Low-Volume, High-intensity Interval Training in Patients with Coronary Artery Disease. *Medicine and science in sports and exercise*,
- [23] K. D. Currie, L. M. Rosen, P. J. Millar, R. S. McKelvie, and M. J. MacDonald, "Heart rate recovery and heart rate variability are unchanged in patients with coronary artery disease following 12 weeks of highintensity interval and moderate-intensity endurance exercise training," *Applied Physiology, Nutrition, and Metabolism*, vol. 38, no. 6, pp. 644–650, 2013.
- [24] G. G. Cardozo, R. B. Oliveira, and P. T. V. Farinatti, "Effects of high intensity interval versus moderate continuous training on markers of ventilatory and cardiac efficiency in coronary heart disease patients," *The Scientific World Journal*, vol. 2015, Article ID 192479, 8 pages, 2015.
- [25] T. Guiraud, A. Nigam, V. Gremeaux, P. Meyer, M. Juneau, and L. Bosquet, "High-intensity interval training in cardiac rehabilitation," *Sports Medicine*, vol. 42, no. 7, pp. 587–605, 2012.
- [26] T. T. Moholdt, B. H. Amundsen, L. A. Rustad et al., "Aerobic interval training versus continuous moderate exercise after coronary artery bypass surgery: a randomized study of cardiovascular effects and quality of life," *American Heart Journal*, vol. 158, no. 6, pp. 1031–1037, 2009.
- [27] T. Moholdt, I. L. Aamot, I. Granøien et al., "Aerobic interval training increases peak oxygen uptake more than usual care exercise training in myocardial infarction patients: a randomized controlled study," *Clinical Rehabilitation*, vol. 26, no. 1, pp. 33–44, 2012.
- [28] T. Moholdt, E. Madssen, Ø. Rognmo, and I. L. Aamot, "The higher the better? Interval training intensity in coronary heart disease," *Journal of Science and Medicine in Sport*, vol. 17, no. 5, pp. 506–510, 2014.
- [29] Ø. Rognmo, E. Hetland, J. Helgerud, J. Hoff, and S. A. Slørdahl, "High intensity aerobic interval exercise is superior to moderate intensity exercise for increasing aerobic capacity in patients with coronary artery disease," *European Journal of Cardiovascular Prevention and Rehabilitation*, vol. 11, no. 3, pp. 216–222, 2004.
- [30] D. E. R. Warburton, D. C. McKenzie, M. J. Haykowsky et al., "Effectiveness of high-intensity interval training for the rehabilitation of patients with coronary artery disease," *American Journal of Cardiology*, vol. 95, no. 9, pp. 1080–1084, 2005.

- [31] U. Wisløff, A. Støylen, J. P. Loennechen et al., "Superior cardiovascular effect of aerobic interval training versus moderate continuous training in heart failure patients: a randomized study," *Circulation*, vol. 115, no. 24, pp. 3086–3094, 2007.
- [32] J. A. C. Sterne, M. Egger, and D. Moher, "Addressing reporting biases," in *Cochrane Handbook for Systematic Reviews of Interventions, Chapter 10*, J. P. T. Higgins and S. Green, Eds., 2011.