

Erratum: Long-term Visual and Refractive Outcomes of Argon Laser-treated Retinopathy of Prematurity

Majid Farvardin^{1,2}, MD; Zahra Kalantari^{1,2}, MD; Mohammadreza Talebnejad^{1,2}, MD; Marzieh Alamolhoda², PhD; Amir Norouzpour³, MD

J Ophthalmic Vis Res 2022; 17 (4): 609–610

In the article titled “**Long-term Visual and Refractive Outcomes of ArgonLaser-treated Retinopathy of Prematurity**” published on pages 384–389, Volume 17, Issue 3 of *Journal of Ophthalmic and Vision Research*,^[1] Table 2 and Figure 1 were missing in the full-text PDF and HTML of the article. The online version of the article has therefore been updated on November 24, 2022 and can be accessed from <https://knepublishing.com/index.php/JOVR/article/view/11576>.

Table 2. Visual, structural, and refractive outcomes of the four groups.

Variables	Laser-treated ROP (Group I) (n = 12)	Spontaneously-Regressed ROP (Group II) (n = 93)	No ROP (Group III) (n = 37)	Control (Group IV) (n = 143)	P value
Corrected distance VA, logMAR (mean ± SD)					
OD	0.09 ± 0.29	0.02 ± 0.14	0.00 ± 0.00	0.01 ± 0.05	0.04*
OS	0.00 ± 0.00	0.01 ± 0.06	0.00 ± 0.00	0.01 ± 0.05	0.14*
Macular dragging or scar					
OD					
No, n (%)	10 (83.3)	92 (98.9)	37 (100)	142 (99.3)	0.01 ^{△△}
Yes, n (%)	2 (16.7)	1 (1.1)	0 (0)	1 (0.7)	
OS					
No, n (%)	11 (91.7)	92 (98.9)	37 (100)	142 (99.3)	0.24 ^{△△}
Yes, n (%)	1 (8.3)	1 (1.1)	0 (0)	1 (0.7)	
Strabismus, n (%)					
No (< 8 PD)	9 (75)	87 (93.5)	36 (97.3)	135 (94.4)	0.04 [△]
Yes (≥ 8 PD)	3 (25)	6 (6.5)	1 (2.7)	8 (5.6)	
Cylindrical power (D) (mean ± SD)					
OD	-1.06 ± 0.32	-0.77 ± 0.50	-0.80 ± 0.66	-0.70 ± 0.87	0.001*
OS	-0.79 ± 0.44	-0.77 ± 0.56	-0.84 ± 0.77	-0.65 ± 0.79	0.004*
Spherical equivalent (D) (mean ± SD)					
OD	0.24 ± 1.95	0.92 ± 1.04	1.04 ± 0.65	0.21 ± 1.64	<0.001*
OS	0.94 ± 1.07	0.86 ± 1.28	1.08 ± 0.77	0.27 ± 1.61	<0.001*
OD					
Myopia (≤ -1 D), n (%)	2 (16.7)	1 (1.1)	0 (0)	15 (11)	
Non-Myopia (>-1 D), n (%)	10 (83.3)	92 (98.9)	37 (100)	121 (89) ⁺	
OS					
Myopia (≤ -1 D), n (%)	1 (8.3)	1 (1.1)	0 (0)	10 (7.4)	
Non-Myopia (>-1 D), n (%)	11 (91.7)	92 (98.9)	37 (100)	126 (92.6) ⁺	

Table 2. (Continued).

Variables	Laser-treated ROP (Group I) (n = 12)	Spontaneously- Regressed ROP (Group II) (n = 93)	No ROP (Group III) (n = 37)	Control (Group IV) (n = 143)	P value
Anisometropia,					
≥ 1.5 D, n (%)	3 (25)	2 (2.2)	1 (2.7)	6 (4.4)	0.03 ^{ΔΔ}
< 1.5 D, n (%)	9 (75)	91 (97.8)	36 (97.3)	130 (95.6) ⁺	

SD, standard deviation; logMAR, the logarithm of the minimum angle of resolution; VA, visual acuity; OD, right eye; OS, left eye; PD, prism diopter; D, diopter

*Kruskal–Wallis; ^ΔChi-square; ^{ΔΔ}Fisher's exact test; ⁺missing data




Figure 1. Ocular biometric parameters among the four groups for the right (OD) and left (OS) eyes. I–IV indicate groups 1 through 4. The first four plots (a–d) are Boxplots analyzed using Kruskal–Wallis test, and the last two plots (e–f) were analyzed using ANOVA. Brackets with an asterisk (*) represent significant differences ($P < 0.05$). D, diopter; mm, millimeter; AC Depth, anterior chamber depth; SEM, standard error of mean.

REFERENCES

- Farvardin M, Kalantari Z, Talebnejad M, Alamolhoda M, Norouzpour A. Long-term visual and refractive outcomes of argon laser-treated retinopathy of prematurity. *J Ophthalmic Vis Res* 2022;17:384–389.