



SEX DETERMINATION OF BALI STARLING (*Leucopsar rothschildi*) USING MOLECULAR SEXING

Putu Indra Pramana Wirastika¹, Ignatius Pramana Yuda², Felicia Zahida
Fakultas Teknobiologi Universitas Atma Jaya Yogyakarta,
Jl. Babarsari 44 Yogyakarta 55281, Indonesia
Email: 1) Nd_ra_21@yahoo.com; 2) pramyd@staff.uajy.ac.id

ABSTRACT

Bali Starling (*Leucopsar rothschildi*) are monomorphic at the age of nestling. For the conservation of bird it important is to determine its sex at the earlier stage. Conventional methods have limitations. This study applied PCR-based molecular sexing to answer this issue. This study aimed to obtain the most effective molecular primers to identify the sex of Bali starling. The most common used combination of P2/P8, 2550F/2718R and 1237L/1272H primers, which amplify CHD1 gene (Chromo-helicase-DNA-binding) were evaluated. DNA samples were obtained from secondary wing feathers of young Bali Starling. Separation in agarose gel electrophoresis of PCR products showed that the three primers were successfully amplified the samples with different degrees of success, that was 90% (P2/P8), 86.7% (2550F/2718R), and 73.3% (1237L/1272H), respectively. However, only the combination of P2/P4 and 2550F/2718R primers was able to sex Bali Starling based on observation of PCR products on agarose gel. The sizes of the genes were slightly different with those reported on previous studies. Most of the results of molecular sexing were in accordance with the sex based on morphological characters.

Key words : Bali starling, *Leucopsar rothschildi*, molecular sexing, CHD gene

INTRODUCTION

Bali Starling is an endemic bird to Bali Island, and due to its small population it has been classified by International Union for Conservation of Nature (IUCN) as Critically Endangered species (BirdLife International, 2013). A breeding program in Tegal Bunder has been initiated to increase the remaining population. Bali Starling is considered sexually monomorphic, particularly in young bird, so it is hard to determine an individual's sex without having the bird in hand. For nestling, however, it is very difficult to identify the sex by morphological examination.

The identification of sex is very important because it is one of keys points on Bali Starling breeding program. By using the knowledge of sex identification genes, the breeding program can be applied more successfully.

Some methods have been developed for sex identification of monomorphic birds, i.e., vent sexing, laparoscopy, sexing steroid and karyotyping. These methods, however, are less or unreliable, time-consuming and expensive. Even some of the methods can hurt or even kill the bird (Dubiec and M, 2006). The DNA-based molecular sexing provides a solution.

The sex of an individual is recognized from the genes located on sex chromosomes. Female birds have one copy of both Z and W, and male birds have two copies of Z avian sex chromosomes. There are many DNA techniques available, such as Griffiths *et al.* (1998) who amplified sex specific CHD1 gene by using P2 - P8 primer pairs, which was successful for 27 of the 28 tested bird species from across the class of Aves. Furthermore Fridolfsson and Ellegren (1999) used 2550F/2718R primer pairs to test the sex of non-ratite birds. This study tested 50 birds species from 11 orders and successfully sexing 47 of the species.

Meanwhile 1237L/1272H primer pairs was used by Kahn *et al.*(1998). Those three combinations of primers are the most common primer sets which are used for molecular sexing. This study aims to assess the reliability of those tree primer set to identify the sex of Bali Starling.

MATERIALS AND METHODS

Samples

Materials for DNA samples were obtained from molted wing feathers (score 2), which were plucked from the young birds of breeding program in Bali Barat National Park. In total 30 samples were obtained from different individual young birds, which were sexually known based on its morphology were examined in this study. Each feather sample was stored in an envelope and then transferred into a 1.5 ml centrifuge tube filled with *Queen's Lysis buffer* until sink, and then it was kept on the room temperature or refrigerator.

DNA extraction was done using terminal part of bird's feather (0,5-1 cm) then it was put into a centrifuge tube containing 500 ml lysis buffer (50mM Tris-HCL pH 8; 20 mM *ethylene diamine tetra acetic* [EDTA], 2% *sodium dodecyl sulfate* [SDS]) and *proteinase K* with consistency amount of 175 mg/ml. Afterwards, samples were incubated at 37°C for 24 hours. After 24 hours the temperature was increased to 50°C for 1 hour. After the lysis process, sample was centrifuged on 13000 rpm for 10 minutes. DNA purification was done by phenol chloroform isoamylalcohol (PCI) with comparison of 25:24:1 (Bello, 2001).

PCR analysis

PCR amplifications were performed in Veriti® thermal cycler, in a 25 iL volume. Each reaction using approximately 20 ng of genomic DNA as template, primers P2/P8 at 0,3 mM each, 0,2 mM each dNTP, 1,5 mM MgCl, 3U Tag DNA polymerase, and 1x Tag Buffer. Meanwhile for the other two primers sets (2550F/2718R and 1237L/1272H), each reaction contained of primers at 0,2 mM each, 0,2 mM each dNTP, 1,5 mM MgCl, 2U Tag DNA polymerase, and 1x Tag Buffer. After a series of optimizing the reaction was reached, the PCR program used for the P2/P8 primers was 94°C for 5 minutes, then 40 cycles of 94°C for 30 sec., 48°C for 45 sec., 72°C for 30 sec, followed by 72°C for 5 min. Meanwhile for the 2500F/2718R primers the PCR conditions was 95°C for 5 minutes, then 35 cycles of 95°C for 30 sec., 42°C for 40 sec., 72°C for 30 sec, followed by 72°C for 5 min; and for 1237L/1272H primers was 92°C for 2 minutes, then 30 cycles of 95°C for 30 sec., 57°C for 45 sec., 72°C for 45 sec, followed by 72°C for 5 min.

DNA fragments obtained by PCR were separated in 3% gel agarose stained with SYBER®Safe, and was examined visually under UV light in KODAK Gel Logic 100 System.

RESULTS AND DISCUSSION

The band patterns of DNA fragment obtained by PCR for Bali Starling were showed in Figure 1. Female samples presented two bands, meanwhile male samples presented one band and smaller size, either for PCR using primer set 2550F/2718R (Figure1A) or P2/P8 (Figure 1C). However, the use of combination primer 1237L/1272H was not able to differentiate the male and female samples. All the samples which were amplified using the primer

set 1237/1272H showed single band. Increasing the concentration of the agarose gel (4%) showed the same results. This finding suggested the need to separate the amplicon on acrylimide gel. The same results were reported by Kahn *et al.* (1998) for seven bird species, that the female bird samples showed single band in agarose gel. However, two of them resolved two bands in acrylimide gel.

Amplification using primer set 2550F/2718R obtained the amplicon size of chromosome Z of 679 bp and the second was chromosome W as 459 bp (Figure 1A). These sizes were slightly different compare to the finding reported by Dawson *et al.* (2001), using the primer to sex four species of auklet (Z=600; W=430). In other had using primer set P2/P8 obtained amplicon size from chromosome W = 393 bp and chromosomes Z =351 bp (Figure 1C). These sizes were in agreement with the study by Griffiths *et al.* (1998). The single band of the fragment DNA which were amplified using the combination 1237L/1272H primers was ~338 bp (Figure 1B) was longer than the range size reported by Kahn *et al.* (1998).




Figure 1. PCR amplifications of three different genotypes using the combination of (A) 2550F/2718R, (B) 1237L/1272H, and (C) P2/P8 primers, run and imaged on a 3% gel and stained with SYBR Safe.

The PCR product suggested that the three common combinations of primers for molecular sexing have successfully amplified the chromosome sex of the Bali Starling. However, among the 30 samples assessed were not all been amplified. The success ratio were 90%, 86,7%, and 73,3 %, respectively for the combination P2/P8, 2550F/2718R and 1237L/1272H primers. Based on these ratios and the ability to differ the sex, male and female, the combination of P2/P8 primers was the most reliable primer to identify the sex of Bali Starling.

The sex of Bali Starling, which were tested using molecular test, either using combination of 2550F/2718R or P2/P8 primers, were identical for the positive samples (Table 1). However, not all of the results were in accordance with the sex determination based on its morphology. Six samples were identified differently, suggested that there was possibility a mistake on sexing young Bali Starling based on its morphological characters.

Table 1. The sex of Bali Starling based on morphological assessment and molecular test

No	Sample code	Bird sex based on		
		Morphology	Molecular	
			P2/P8	2550F/2718R.
1.	TNBB 0483	♀	♂	♂
2.	TNBB 0488	♂	+	-
3.	TNBB 0061	♂	+	+
4.	TNBB 0484	♂	♂	♂
5.	TNBB 0302	♂	♂	♂
6.	TNBB 0134	♂	♂	♂
7.	TNBB 0029	+	+	+
8.	TNBB 0900	♂	♂	♂
9.	-	+	♂	♂
10.	TNBB 0037	+	♂	♂
11.	TNBB 0056	+	+	+
12.	TNBB 0296	♂	♂	♂
13.	TNBB 0323	+	+	+
14.	TNBB 0336	♂	♂	♂
15.	TNBB 0381	+	+	-
16.	TNBB 0107	♂	♂	♂
17.	TNBB 0398	♂	♂	♂
18.	TNBB 0146	+	+	+
19.	TNBB 0337	+	+	+
20.	TNBB 0368	♂	♂	♂
21.	TNBB 0131	+	♂	♂
22.	TNBB 0301	+	+	+
23.	TNBB 0415	♂	♂	♂
24.	TNBB 0405	+	+	+
25.	TNBB 0211	♂	♂	♂
26.	TNBB 0491	+	+	-
27.	-	+	+	+
28.	TNBB 0399	♂	-	♂
29.	TNBB 0419	♂	-	♂
30.	TNBB 0420	+	-	+

Note: ♀ = female, ♂ = male

ACKNOWLEDGMENTS

We thanks to the Head of Bali Barat National Parks for the permission for DNA sampling. This research received fund from DIKTI.

REFERENCES

- BirdLife International. 2013. Species factsheet: *Leucopsar rothschildi*. .
- Dawson, D. A., S. Darby, F. M. Hunter, and A. P. Krupa. 2001. A critique of avian CHD-based molecular sexing protocols illustrated by a Z-chromosome polymorphism detected in auklets. *Molecular Ecology Notes* 1:201-204.
- Dubiec, A. and Z.-N. M. 2006. Molecular Techniques For Sex Identification In Birds. *Biological Letters* 2006 43 (1): 3.12. 43:3-12.
- Fridolfsson, A.K., and H. Ellegren. 1999. A simple and universal method for molecular sexing of non-ratite birds. *Journal of Avian Biology* 30:116-121.
- Griffiths, R., M. C. Double, K. Orr, and R.J.G. Dawson. 1998. A DNA test to sex most birds. *Molecular Ecology* 7:1071-1075.

Kahn, N. W., J. S. John, and T. W. Quinn. 1998. Chromosome-Specific Intron Size Differences in the Avian CHD Gene Provide an Efficient Method for Sex Identification in Birds. *The Auk* 115:1074-1078.