

Conference Paper

Rare Color Variants in Lacertidae on the Example of *Zootoca vivipara* (Jacquin, 1787) in Ukraine

O.D. Nekrasova

I. I. Schmalhausen Institute of Zoology of National Academy of Sciences of Ukraine, st. Bogdana Khmel'nitskogo, 15, Kyiv, 01030, Ukraine

Abstract

We used GIS modeling (DIVA-GIS, Maxent) to simulate the potential distribution of *Zootoca vivipara*. For modeling, we used a database including information from our studies and literature, comprising 190 points of lizard samples (including 7 points for melanistic lizards). It can be concluded that the distribution of *Z. vivipara* in Ukraine depends on the following environmental variables related to solar radiation and precipitation: "Radiation of wettest quarter" (BIO24; 34.2%), "Precipitation of driest week" (BIO14; 18.1%), "Lowest weekly radiation" (BIO22; 15.7%) and "Mean diurnal temperature range" (BIO2; 12.8%). For habitats where melanistic lizards were found, "Mean moisture index of coldest quarter" (BIO35; 18.9%) was also significant in addition to BIO24 (50.4%) and BIO22 (17.3%).

Keywords: melanistic lizards, *Zootoca vivipara*, GIS modeling, Ukraine.

Corresponding Author:

O.D. Nekrasova
 oneks22@gmail.com

Received: 23 January 2018

Accepted: 20 April 2018

Published: 3 May 2018

Publishing services provided by
 Knowledge E

© O.D. Nekrasova. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Selection and Peer-review under the responsibility of the Amphibian and Reptiles Anomalies and Pathology Conference Committee.

1. Introduction

In the literature, there are many descriptions of the rare color variants of lizards, such as morphological aberrations (anomalies) and morphs. Abnormalities in reptiles and amphibians are the result of the disturbed function of chromatophores and other elements which form the coloring [24]. For example, melanophores (black and brown), xanthophores (yellow), erythrophores (red), iridophores (iridescent, blue, silver or gold), cyanophores (blue) and leucophores (dull, whitish) are widespread [8]. Reptiles exhibit varying chromism, which is defined by partial or complete absence of different types of pigment cells. Melanism, the opposite of albinism, is the development of a dark-colored melanin pigment in the skin. The presence, absence, density and distribution of melanocytes and chromatophores within each layer determine the color of the reptile: for example, the color variants "nigra" or "platini" in *Lacerta agilis* Linnaeus, 1758 [19]. A mosaic of combined black and brown colors is described in the literature as "brown melanism" [25, 26]. Pseudo-melanism, also called abundism, is

OPEN ACCESS

another variant of pigmentation, one characterized by dark spots or enlarged stripes which cover a large part of animal's body, making it look melanistic. For some reptilian species, a completely black body is normal and is maintained by natural selection, for example, *Vipera (Pelias) nikolskii* Vedmederja, Grubant et Rudaeva, 1986 (*V. berus nikolskii*). The frequency of the occurrence of rare color variants among reptiles depends on the uniqueness of their population. If the black color version is common, then the morph (or variant) is written as "var. nigra" [10] (more than 5%). Isolated cases of deviations from the normal state are called aberrations – "aber. nigra" – from the normal color (less than 5%). For example, in Ukraine such aberrations were found among melanistic *L. agilis* on Island Biruchiy [33, 39] and in the Molochnij estuary [15]; among specimens of *Lacerta viridis* (Laurenti, 1768), they were found in Mykolaiv region [23]; among *Podarcis taurica* (Pallas, [1814]) in Crimea [14]; among *Darevskia lindholmi* (Lantz et Cyrén, 1936) in Crimea [19]; and among *Darevskia dahlia* (Darevsky, 1957) ("brown melanism") in the Zhytomyr region [25]. Melanistic morphs were described among *L. agilis* and *Zootoca vivipara* (Jacquin, 1787) in Russia. Singular melanistic specimens of *L. agilis* were found in Krasnodar krai [35], Western Altai [19, 39] and the Stavropol krai of Samara province [7]. Melanistic morphs have been found among *Z. vivipara* more often than in other reptile species in Russia. In Sverdlovsk province, the frequency of melanistic individuals in populations in forests was 1.15%, and in countryside it ranged from 0.95% to 1.1% [37]. Up to 5.5% of specimens were melanistic (more among the females more) in the Cherdyn district of Perm province [7]. Melanistic individuals are also reported in Leningrad province [21] and near Tomsk [20]. In Europe (the United Kingdom, Spain, Germany, Ireland, Switzerland, Poland, Slovakia, Montenegro, etc.), melanistic specimens are more common in *Z. vivipara* than in other species of the Lacertidae family [1, 10, 12, 27, 28, 38]. In the Czech Republic, up to 8% of *Z. vivipara* specimens were melanistic, and this variant of coloring is much more likely to be found among males [11]. In Denmark, 10% of lizards were melanistic [9]. Earlier in the literature, in 1805, black variant lizards (*Lacerta nigra*) were described in Wolf in Sturm (the type locality is "Schneegebirge, so genannte Wenger Alpe", in the canton of Berne, Switzerland; Schmidtler, Böhme, 2011). In July 1850, the *L. nigra* ("*L. Fitzingeri*") was found near Kharkov (Ukraine; [3]). Chromium is known for *Z. vivipara* (male) to be green in the UK (Bowles, 2003; [5, 37]). Rare cases of melanism in the Balkan Peninsula, near the southern range limits for this species, were described in the literature [12]. Studies of rare abnormalities (or variants) in the coloration of lizards have not been conducted in Ukraine yet. Therefore, the aim of our work is to compile

information about the cases of melanistic *Z. vivipara* in Ukraine in order to identify the potential bioclimatic factors that could lead to such color manifestations.

2. Methods

To explore this issue, we used GIS modeling of the potential distribution of *Z. vivipara* revealed in the territory of Ukraine (183 points and 7 points of melanistic findings). *Z. vivipara* var. *nigra* were found in regions of Ukraine like: Ivano-Frankivsk (Polonyna Bryusny; reported Zelenchuk Ya., Smirnov N., 2015; Fig. 1: №1; 47.84° N, 24.74° E), Sumy (Desnyansko-Starogutsky NNP, near the village of Ochkino according to Kotserzhinskaya, 2003; Fig. 1: №2; 52.24° N, 33.38° E), Kyiv (the village of Pluty; Fig. 1: №3; 50.16° N, 30.71° E), Cherkassy (Sytnik, 2004; Fig. 1: №4; 49.72° N, 31.49° E), Poltava - NNP "Pyryatynsky" (Udaj River, 2014; Fig. 1: №5; 50.23° N, 32.53° E), around the village of Kamyshnaya (Khorol River, 2012; Fig. 1: №6; 50.18° N, 33.72° E) and Kharkiv (NNP "Slobozhansky"; reported Biatov A.; Fig. 1: №7; 50.05° N, 35.18° E).




Figure 1: The results of GIS modeling of potential distribution of *Z. vivipara* (DIVA-GIS and Maxent) (No. 1-7 marking in the text).

These data were collected from the results of original research (findings, verified information) and the literature [16, 29, 40]. They were the basis for a database. Mapping, modeling and calculations were carried out using OziExplorer v.3.95.4 m, DIVA-GIS v.5.2. (19 bioclimatic factors from WorldClim database, Domain algorithm) and Maxent v. 3.3.3k. (35 bioclimatic factors) [36].



Figure 2: Melanistic viviparous lizard from the Poltava region.

3. Results

Viviparous lizards are widely distributed throughout the whole Eurasian forest zone. Its range extends to the north of the Arctic Circle, so it is tolerant of low temperatures. These lizards may appear very early in spring, before the snow melts away (in March, when the air temperature is around +4°C). The southern range limit runs at 48° northern latitude. In these areas, viviparous lizards live in damp locations, often near water (along rivers and in swamps and damp forests). The background coloration among these lizards is very variable - most animals are basically brown (chocolate colored), and grey or olive is less likely. In contrast to the relatively low morphological variability characteristic of this species, there is usually a high interpopulation variation in patterns [1]. Juveniles start by being black or chocolate colored, and slowly acquire a pale green ventral coloration on becoming yearlings. We recorded melanistic *Z. vivipara* (young-adults) in 7 localities (Fig. 1., Fig. 2.). In the Poltava region near the Khorol River, only one lizard specimen was black (n=15; Fig. 1: N°6). Most of the finds were singular. Along River Khorol, 3-5 to 10 lizards per km were found.

GIS modeling showed that the distribution of *Z. vivipara* in Ukraine depends on the following parameters related to solar radiation and precipitation: "Radiation of wettest quarter" (BIO24; 34.2%), "Precipitation of driest week" (BIO14; 18.1%), "Lowest weekly radiation" (BIO22; 15.7%) and "Mean diurnal temperature range" (BIO2; 12.8%). For habitats where melanistic lizards were found, "Mean moisture index of coldest quarter" (BIO35; 18.9%) was also significant in addition to BIO24 (50.4%) and BIO22 (17.3%). The model presented a high fit - AUC = 0.996. Functions of coloration

may be very different, for example: camouflage, UV protection, behavioral adjustments, ecology, thermobiology, etc. Melanism is thought to have thermoregulatory significance and is considered to be a result of ecological adaptation to a certain environment [4, 12, 34]. Also, some scholars have opined about the thermobiological benefits of a melanistic (dark-colored) reptile, arguing that such animals heat up faster, which allows them to be more active in adverse conditions. The skin reflectivity of melanistic wall lizards was 2% higher than that of non-melanistic ones in a short wavelength range 0.38-0.80 μ [34]. There are different opinions about “the thermal melanism hypothesis (TMH)” and its specificity with regards to various regions [4]. Exploring some thermobiological features of melanistic *Z. vivipara* (heating rate, body size and habitat) in the Czech Republic, researchers could not confirm the expected benefit of melanistic color. It was suggested that the occurrence of melanistic specimens among viviparous lizards is due to the characteristics of the habitat (vegetation, background color, habitat, etc.) [11]. So, the manifestation of melanism and changes in the color and body shape in different reptilian species is also associated with a complex of factors – extreme and edge effects, height above sea level, composition of the soil (e.g., excessive quantities of substances that stimulate melanin production: manganese and iron), etc. [6, 30]. For example, lizards inhabiting more humid, dark areas are darker than species in dry open habitats [37]. Also, the light color of *Vipera (Pelias) berus* (Linnaeus, 1758) has more xerophiles and thermophiles (forest edges, southern slopes) than black snakes (river banks or swamp edges) [22].

4. Conclusion

It can be concluded that black skin color can be influenced by a complex of factors (physical, chemical, ecological, marginal, biological and climatic). In some cases, the occurrence of rare colors in lizards may be seen as an adaptation to a certain environment. Each case should be considered individually and regionally. We also observed the possible influence of climatic features on the occurrence of melanistic forms among viviparous lizards: in addition to factors associated with solar radiation and precipitation (“Radiation of wettest quarter” (BIO24; 34.2%) and “Lowest weekly radiation” (BIO22; 15.7%)), “Mean moisture index of coldest quarter” (BIO35; 18.9%) is also significant. In fact, all such specimens were discovered near the southern border of the area (Fig. 1: №1, 3-7): in similar habitats, there are live black reptile species, such as *V. nikolskii*. Therefore, we can assume that the expression of melanism is not accidental and not subject to selection. Since data on the occurrence of melanistic forms among

viviparous lizards is not adequate, research should continue. However, according to preliminary data, it is important to note that in some regions where melanistic specimens are quite common (over 5%), especially among young individuals, this manifestation can be considered a variation on the normal color of the viviparous lizard of *Z. vivipara* var. *nigra* [10].

Acknowledgements

The author is thankful to Marushchak A., Shevchenko O. and Oskyrko O. for their assistance.

References

- [1] Boulenger GA: On the variation of the common lizard, *Lacerta vivipara*. Journal of Zoology Research. 1917; 2: 1-16.
- [2] Bowles FD: *Zootoca vivipara* (Common or Viviparous lizard): green colouration. Herpetol. Bull. 2000; 85: 29.
- [3] Chernay AV: Fauna Khar'kovskoy gubernii i prilozhashchikh k ney mest. Fauna zemnovodnykh zivotnykh i ryb. Akt v imp. Khar'kov. un-te, 1850; V. 1: 5-6.
- [4] Clusella-Trullas S, Terblanche JS, Blackburn TM, & Chown SL: Testing the thermal melanism hypothesis: a macrophysiological approach. Functional ecology. 2008; 22(2): 232-238.
- [5] Palmer K: *Lacerta (Zootoca) vivipara* (Common or Viviparous lizard): alternate green colour phase. Herpetol. Bull. 2005; 92: 30.
- [6] Doronin IV: Cases of melanism in lizards of the genus *Darevskia*. Zoologicheskyy zhurnal; 91.11. 2012: 1420-1427 [in Russian].
- [7] Eplanova GV: The findings of melanistic specimens of *Lacerta agilis* and *Zootoca vivipara*. In: Proceedings of the 5th Congress of the A.M. Nikolsky Herpetological Society: "The Problems of Herpetology" (24-27 September 2012, Minsk) Belarus; 2012: 84-87 [in Russian].
- [8] Fujii R: The regulation of motile activity in fish chromatophores. Pigment Cell Research. 2000; 13(5): 300-319.
- [9] Glandt D: Die Waldeidechse: unscheinbar - anpassungsfähig - erfolgreich. Z. Feldherpetologie, Beiheft 2. Laurenti-Verlag, Bochum; 2001: 111.
- [10] Gray JE: Note on *Zootoca vivipara* var. *nigra*. Ann. Mag. Nat. Hist. 1857; 19 (2): 107.

- [11] Gvoždík L: Colour polymorphism in a population of the common lizard, *Zootoca vivipara* (Squamata: Lacertidae). *Folia Zoologica*. 1999; 48: 131-136.
- [12] Iković V, & Gvozdrenović S: A record of melanistic viviparous lizard *Zootoca vivipara* (Lichtenstein, 1823) (Squamata, Lacertidae) on Prokletije Mountain, Montenegro. *HYLA-Herpetološki bilten*. 2014; 2: 41-44.
- [13] Jambrich A, & Jandzik D: Melanism in the topotypic population of the Pannonian subspecies of the common lizard, *Zootoca vivipara pannonica* (Reptilia: Lacertidae). *Herpetology Notes*. 2012; 5: 219-221.
- [14] Karmishev YuV: Find mechanistically specimen of the *Padarcis taurica* in the Steppe Crimea. *Vestnik zoologii*. 2001; 35(1): 58 [in Russian].
- [15] Karmishev YuV: Modern state of reptiles fauna coastal area of Molochnij estuary. *Issues of Ecology and Bioindication*. Zaporozhye, ZNU, 2014; 19 (1): 199-210 [in Russian].
- [16] Korshunov AV, Zinenko AS: Features of the biotopic distribution reportaway lizards *Lacerta vivipara* on the periphery of the range (Kharkov region). In: *Proceedings of the II international scientific conference "Biodiversity and role of zoocenosis in natural and anthropogenic ecosystems: Zoocenosis – 2003"*. Dnepropetrovsk: DNU; 2003: 206-208 [in Russian].
- [17] Kotenko TI, Sviridenko YeYu: Variability of coloration and pattern of the sand lizard, *Lacerta agilis* (reptilia, sauria, lacertidae): methodological aspects. *Vestnik zoologii*. 2010; 44(2): 137-162 [in Russian].
- [18] Kotserzhinskaya IM: The herpetofauna desniansko-Starogutsky national natural park and adjacent areas. *Zapovidna sprava v Ukraini*. 2003; 9(2): 45-51 [in Russian].
- [19] Kukushkin OV, Sviridenko EYu: Finds of melanistic specimens of a rock lizard *Darevskia lindholmi* (reptilia, sauria, lacertidae) in the Crimea. *Vestnik zoologii*. 2002; 36(3): 98 [in Russian].
- [20] Kuranova VN: About melanism *Zootoca vivipara* and *Vipera berus*. *Vestnik zoologii*. 1989; 2: 59-61.
- [21] Maymin MYu, Orlov NL: Three cases of melanism have reptiles of Squamata. In: *Proceedings of the "problems of herpetology"*. Leningrad: Nauka;1977: 141 [in Russian].
- [22] Litvinov NA, Ganshchuk SV: Thermodaptations of the reptiles of the Volga basin. *Izvestiya Samarskogo scientific center, Russian Academy of Sciences*. 2010; 12(1): 133-137 [in Russian].
- [23] Loparev SA, Tzvelych AN: Finding of melanistic individuum of *Lacerta viridis* Laur. *Vestnik zoologii*. 1981; 4: 75 [in Russian].

- [24] Nekrasova OD: Some aspects of anomaly manifestation in amphibian coloration. In: Proceeding of the international conference "Anomalies and Pathologies of Amphibians and Reptiles: Methodology, Evolutionary Impact, Possibilities for Estimation of Environmental Health" (23-26 September 2013, Yekaterinburg, Russia): 144-149 [in Russian].
- [25] Nekrasova OD, Kostiusyn VA: Current distribution of the introduced rock lizards of the *Darevskia (Saxicola)* complex (Sauria, Lacertidae, Darevskia) in Zhytomyr region (Ukraine). *Vestnik zoologii*. 2016; 50(3): 225-230.
- [26] Ostrovskikh SV: Different form of melanism and its development with age in the populations of stepper viper *Vipera renardi* (Christoph, 1861). *Russ. J. Herpetol*. 1997; 4(2): 186 – 191.
- [27] Petzold HG: Nigrinos von *Lacerta vivipara* aus der Umgebung Berlins (Reptilia: Sauria: Lacertidae). *Salamandra*. 1978; 14: 98-100.
- [28] San-Jose LM, Gonzales-Jimena V & Fitze PS: Frequency and phenotypic differences of melanistic and normally coloured common lizards, *Lacerta(Zootoca) vivipara* of the Southern Pyrenees (Spain). *Herpetological review*. 2008; 39: 422-425.
- [29] Shaitan SV: Peculiarities of distribution and ecology of amphibians and reptiles (amphibia, reptilia) of the western forest-steppe of Ukraine. *Vestnik zoologii*. 1999; 33 (4-5): 95-98.
- [30] Sharygin SA, Popov VN: Phenotypic variability and some geochemical ecology of these lizards of the Crimea. *Actual problems of Herpetology and toxinology*. 2003; 6. Togliatti: 110–116 [in Russian].
- [31] Sytnik AI: True lizards (Lacertidae) in conditions of the central forest-steppe of Ukraine. K.: PPU "Kyiv University"; 2004: 125 [in Ukrainian].
- [32] Schmidler JF, Böhme W: Synonymy and nomenclatural history of the common or viviparous lizard, by this time: *Zootoca vivipara* (Lichtenstein 1823). *Bonn Zool. Bul.* 2011; 60(2): 145-159.
- [33] Szczerbak NN: Reptiles as a model to study the changes of animals in the acclimatization process. In: Proceedings of herpetol. conf. "Questions of Herpetology": Leningrad, Publishing LSU; 1964: 78-79 [in Russian].
- [34] Tosini G, Lanza B, Bacci M: Skin reflectance and energy input of melanic and non-melanic populations of wall lizard (*Podarcis muralis*). In: Korsós, Z. & Kiss, I. (eds.). Proceedings of the sixth ordinary general meeting, Budapest (SEH-HNHM); 1991: 443-448.

- [35] Trofimov AG, Tzvelych AN: On the records of melanistic specimens of the sand lizard (*Lacerta agilis*) and slow worm (*Anguis fragilis*). In: Proceedings "Ecology and systematics of amphibians and reptiles". L.; 1979:120-121 [in Russian].
- [36] Tytar VM: Analysis of home ranges in species: an approach based on modeling the ecological niche. Vestnik zoologii. 2011. Suppl. N 25: 96 [in Ukrainian].
- [37] Vershinin VL: Amphibians and reptiles of Urals. Yekaterinburg: [Ipaе UB RAS]; 2007: 169 [in Russian].
- [38] Westrin L: Melanistic common lizard, *Lacerta vivipara* (Jacquin), found in Sweden. Fauna och. flora, 1985; 80: 37-38.
- [39] Yablokov AV, Baranov AS, Rozanov AS: Reconstruction of the species Microphylogenesis (as exemplified by studying *Lacerta agilis* L.). Vestnik zoologii. 1981; 3: 11-16 [in Russian].
- [40] Zabroda SN: Amphibians and reptiles of Central and Western Polissya: PhD thesis: spec. 03.00.08 "Zoology". K., 1983: 16 [in Russian].