

## Research Article

# Pregnancy and Neonatal Outcomes: A Study During the COVID-19 Pandemic

Awatiful Azza<sup>1\*</sup>, Nikmaturrahmah<sup>2</sup>, and Ilanka Cahya Dewi<sup>3</sup><sup>1</sup>Department of Maternity Nursing Faculty of Health Science Universitas Muhammadiyah Jember<sup>2</sup>Department of Pediatric Nursing Faculty of Health Science Universitas Muhammadiyah Jember<sup>3</sup>Faculty of Engineering, Universitas Muhammadiyah Jember**ORCID**Awatiful Azza: <https://orcid.org/0000-0003-0946-1092>**Abstract.**

During the COVID-19 pandemic, there were many large-scale restrictions on routine services, including maternal and neonatal health services. This study used a correlational design to analyze the relationship between pregnancy during COVID-19 and the neonate's condition. Samples are pregnant women between 2020 and 2021, totaling 108 samples obtained by random sampling. Data collection used questionnaires, structured interviews, and documentary studies. The approach used was a retrospective about Ante Natal Care visits during COVID-19 with the birth weight of the baby. Data analysis was performed using the Chi-square. There is a relationship between pregnant women's and babies' weight during COVID-19 with a p-value of 0.016 and an OR of 3.967 (95% CI 1.22-12.879). In addition, the number of respondent visits is also related to babies' birth weight (P-value of 0.004 with OR 168 (95% CI 0.045-0.625). Meanwhile, the size of the mother's upper arm circumference and blood pressure is unrelated to the baby's birth weight. During the COVID-19 period, restrictions on Ante Natal Care visits caused pregnancies to become out of control. Lack of education about nutrition and stress levels of pregnant women contributes to pregnancy outcomes.

**Keywords:** baby's birth weight, covid 19 period, pregnant women

## 1. Introduction

Pregnancy is a sensitive period for women in their life cycle. Hormonal changes resulting from adaptation to fetal growth and development in the womb result in physical and psychological changes (1). Physical and psychological changes during pregnancy can be a stressor that causes anxiety in pregnant women (2). Every pregnant woman must receive ante-natal care services according to WHO protocol recommendations, and this is important so that complications during pregnancy can be identified early (3).

During the COVID-19 pandemic, there were many large-scale restrictions on almost all routine services, including maternal and newborn health services. WHO has also set rules regarding antenatal care (ANC) services for pregnant women (4,5). As a

Corresponding Author: Awatiful  
Azza; email: awati-  
ful.azza@unmuhjember.ac.id

Published 8 March 2023

Publishing services provided by  
Knowledge E

© Awatiful Azza et al. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Selection and Peer-review under the responsibility of the ICMEDH Conference Committee.

**OPEN ACCESS**

group was vulnerable to transmission of COVID-19, mothers are not recommended to frequent ANC visits to health services (2,3). Mothers are expected to be able to independently identify complaints and abnormalities that occur in their pregnancy (5). This condition causes the development and health of the mother and baby cannot to be monitored optimally. Mothers and babies are significantly at risk of experiencing pregnancy complications and growth disorders which will impact the outcome of the mother's subsequent pregnancy (4,6).

Data on the impact of pregnant women with COVID-19 in US medical centers for 2020-2021 report that COVID-19 has increased mortality, the need for intubation and ventilation, and admission to intensive care units by 2.1%. Meanwhile, a study in Canada stated that pregnant women during the COVID-19 period experienced a 21.4% tendency to be obese, required 20% intensive care, and experienced premature labor by 12.2% (4).

The Minister of Health of the Republic of Indonesia Number 43 of 2016 Article 2 concerning Minimum Service Standards in the Health Sector states that the statement of integrated antenatal care standards is "Every pregnant woman gets antenatal care according to standards (7,8). WHO protocol policies strongly influence ANC services during the Corona Virus Disease 2019 (Covid19) pandemic. The protocol regulates how pregnant women access health facilities during a pandemic based on existing protocols and maximizes telemedicine use. In addition, the concern about pregnant women being exposed to COVID-19 and the call to stay at home resulted in decreased visits to health facilities (7). This study aims to analyze pregnancy and neonatal outcomes during the COVID-19 pandemic.

## 2. Materials and Methods

This study uses a retrospective cross-sectional approach to pregnant women during the COVID-19 pandemic. The population is all pregnant women in the I, II, and III trimesters recorded in the April 2020-December 2021 period; 108 pregnant women were involved in this study using the random sampling technique in East Java. They were collecting data using a structured questionnaire that has been tested for validity and reliability, with variables being the number of visits by pregnant women, arm circumference, mother's weight, blood pressure, and baby's weight. Data analysis was done with chi-square to assess the proportion comparison between the independent and dependent variables.

### 3. Result

Based on demographic data, it was found that the education of most respondents was primary school 43.5%, with the majority of respondents' pregnancy status being pregnant with their first child (88%). When viewed from the ethnic group, most of the respondents are Madurese (77.8%). The blood pressure data found that some respondents were in a risk group of 66.7%.

The study results in table 2 show a relationship between pregnant women's and babies' weight during the covid pandemic with a p-value of 0.016 and an OR of 3.967 (95% CI 1.22-12.879). In addition, the number of respondent visits is also related to babies' birth weight with a P-value of 0.004 with OR 168 (95% CI 0.045-0.625). Meanwhile, the size of the mother's upper arm circumference and blood pressure is unrelated to the baby's birth weight.

TABLE 1: Demographics of pregnant women during the pandemic (n=108).

Variable	Frequency	Percentage
Education Primary school Junior high school Senior high school	47 23 38	43.5 21.5 35.2
Pregnancy status First pregnancy Second pregnancy More than 2	95 9 4	88 8.3 3.7
Ethnicity Javanese Madurese	24 84	22.2 77.8
ANC visit In accordance Not followed	50 50	50 50
Blood pressure No risk Have risk	36 72	33.3 66.7
Conjunctiva Anemic No anemic	85 23	78.7 21.3
Upper arm circumference Normal Less or more	95 13	88 12
Pregnant woman's weight Normal Abnormal	91 17	84.3 15.7
Birth weight Normal Abnormal	91 17	84.3 15.7

### 4. Discussion

The increase in maternal weight during pregnancy can affect the baby's birth weight. Weight gain during pregnancy is a form of body adaptation because a new individual is growing in the mother's womb (9). In the third trimester, approximately 90% of maternal weight gain is used for fetal growth, placenta, and amniotic fluid. In the first trimester,

TABLE 2: The relationship between weight, ANC visits, upper arm circumference, and blood pressure on the weight of newborns during the covid 19 pandemic (n=108).

Variable	birth weight		P value	OR	95 % CI
	low	normal			
<b>A pregnant woman's weight</b>					
Normal	6	11	0,016	3,967	1,22 - 12,879
Abnormal	11	80			
<b>ANC Visit</b>					
in accordance	3	51	0,004	168	0,045 - 0,625
it is not following	14	40			
<b>Upper arm circumference</b>					
Normal	4	9	0,116	2,803	0,75 - 10,43
Less or more	13	82			
<b>Blood pressure</b>					
No risk	3	33	0,213	3,210	0,95 - 11,23
Risk	14	58			

almost entirely is part of the mother, while in the second trimester, as much as 60% is part of the mother (9,10).

The Indonesian Ministry of Health stated that the weight gain of pregnant women usually ranges from 9-12 kg, while fat deposits in the mother's body are around 3-3.5 kg (11). Maternal weight gain during pregnancy positively correlates with the baby's birth weight (9,12). Pregnant women must meet the need for balanced nutrition during pregnancy so that the baby is born healthy. During the COVID-19 pandemic, pregnant women were very limited in accessing health services, and this was because pregnant women were a group that was vulnerable to contracting COVID (5,13). This condition could cause the mother not to be able to know the progress of her pregnancy correctly. Inappropriate eating patterns during the pandemic and activity restrictions cause weight gain during pregnancy and cannot be monitored properly (14). Limiting ANC activities in essential services and hospitals strongly impacts pregnant women (15).

Antenatal Care (ANC) examination is an essential part of the initial screening of the condition of the baby to be born(6). Poor quality of antenatal care can have an impact on fetal growth and development. Examining upper arm circumference and blood pressure during ANC is a screening to determine the risks pregnant women and fetuses face. Upper arm circumference is one of the screenings to measure the nutritional status of pregnant women (16). If it is less than 23.5 cm, the pregnant woman is considered to be chronically lacking in energy, so pregnant women can be at risk of giving birth to

low birth weight (LBW) babies. This study found that some respondents had an average upper arm circumference of 23.5 cm, as many as 88% (5,16).

Another essential examination during ANC is the measurement of blood pressure. Hypertension in pregnant women is one of the causes of maternal and infant mortality (17). This situation occurs because hypertension in pregnant women can damage the blood vascular system, thereby interfering with exchanging oxygen and nutrients through the placenta from the mother to the fetus (18). This can lead to placental prematurity, which results in slow fetal growth. In addition, constricting blood vessels in pregnant women who suffer from hypertension will increase uterine muscle tone (19). The results of this study indicate that 66.7% of respondents' blood pressure during the pandemic period is a risk group. This condition may be reinforced by high social stressors during the pandemic and activity restrictions so that mothers do not receive education from health workers about the health of their pregnancy. Hypertension during pregnancy can cause complications such as hypofibrinogenemia, hemolysis, brain hemorrhage, eye disorders, pulmonary edema, liver necrosis, HELLP syndrome, and kidney disorders which are significantly at risk of increased maternal and infant morbidity and mortality (20–23).

## 5. Conclusion

There is a relationship between the mother's weight and the number of ANC visits with the baby's birth weight. More intensive supervision is needed for pregnant women to identify the mother's health and the fetus's development.

## Acknowledgments

We gratefully acknowledge the funding from LPDP Through Kementerian Pendidikan, Kebudayaan, Riset, dan Teknologi Direktorat Jenderal Pendidikan Tinggi, Riset, dan Teknologi. We Also Thank The LPPM Universitas Muhammadiyah Jember, Kemuning Lor Arjasa Village, and The Faculty Of Health Sciences Universitas Muhammadiyah Jember.

## References

- [1] Meija L, Rezeberga D. Guidelines: Proper maternal nutrition during pregnancy planning and pregnancy: a healthy start in life Recommendations for health care

- specialists - WHO-OMS. WHO. 2017;1–31.
- [2] Rocca-Ihenacho L, Alonso C. Where do women birth during a pandemic? Changing perspectives on Safe Motherhood during the COVID-19 pandemic. *J Glob Health Sci.* 2020;2(1):4–5.
- [3] Wu H, Sun W, Huang X et al. Online Antenatal Care During the COVID-19 Pandemic: Opportunities and Challenges. *J Med Internet Res* 2020;22(7). <https://doi.org/102196/19916>
- [4] Landrian A, Mboya J, Golub G, Moucheraud C, Kepha S, Sudhinaraset M. Effects of the COVID-19 pandemic on antenatal care utilisation in Kenya: a cross-sectional study. *BMJ Open.* 2022 Apr;12(4):e060185.
- [5] Chmielewska B, Barratt I, Townsend R, Kalafat E, van der Meulen J, Gurol-Urganci I, et al. Effects of the COVID-19 pandemic on maternal and perinatal outcomes: a systematic review and meta-analysis. *Lancet Glob Health.* 2021 Jun;9(6):e759–72.
- [6] Rolnik DL, Matheson A, Liu Y, Chu S, Mcgannon C, Mulcahy B, et al. Impact of COVID-19 pandemic restrictions on pregnancy duration and outcome in Melbourne, Australia. *Ultrasound Obstet Gynecol.* 2021 Nov;58(5):677–87.
- [7] Kemenkes RI. Pedoman pelayanan antenatal, persalinan, nifas, dan bayi baru lahir di Era Adaptasi Baru. 2020. 98 pp.
- [8] Kemenkes. Peraturan Menteri Kesehatan Republik Indonesia Nomor 43 Tahun 2016. 2016
- [9] Simeone RM, Downing KF, Wallace B, Galang RR, DeSisto CL, Tong VT, et al. Changes in rates of adverse pregnancy outcomes during the COVID-19 pandemic: a cross-sectional study in the United States, 2019–2020. *J Perinatol.* 2022 May;42(5):617–23.
- [10] Koenig MD. Nutrient Intake During Pregnancy. *J Obstet Gynecol Neonatal Nurs.* 2017;46(1):120–2.
- [11] Kemenkes RI. Permenkes No 43 Tahun 2016 Tentang SPM Bidang Kesehatan. Kemenkes Republik Indones. 2016;5(2719):1–79.
- [12] Warriner S. Pregnancy and body mass index (BMI). *Oxford Univ Hosp;* 2016. p. 50.
- [13] Karimi L, Makvandi S, Vahedian-Azimi A, Sathyapalan T, Sahebkar A. Effect of COVID-19 on Mortality of Pregnant and Postpartum Women: A Systematic Review and Meta-Analysis. *J Pregnancy.* 2021.
- [14] Atkinson L, De Vivo M, Hayes L, Hesketh KR, Mills H, Newham JJ, et al. Encouraging Physical Activity during and after Pregnancy in the COVID-19 Era, and beyond. *Int J Environ Res Public Health.* 2020 Oct;17(19):1–9.
- [15] Chen SC, Lai YH, Tsay SL. Nursing perspectives on the impacts of COVID-19. *J Nurs Res.* 2020 Jun;28(3):e85.

- [16] Allotey J, Stallings E, Bonet M, Yap M, Chatterjee S, Kew T, et al.; for PregCOV-19 Living Systematic Review Consortium. Clinical manifestations, risk factors, and maternal and perinatal outcomes of coronavirus disease 2019 in pregnancy: living systematic review and meta-analysis. *BMJ*. 2020 Sep;370:m3320.
- [17] Muti M, Tshimanga M, Notion GT, Bangure D, Chonzi P. Prevalence of pregnancy induced hypertension and pregnancy outcomes among women seeking maternity services in Harare, Zimbabwe. *BMC Cardiovasc Disord*. 2015 Oct;15(1):111.
- [18] Berhe AK, Ilesanmi AO, Aimakhu CO, Mulugeta A. Effect of pregnancy induced hypertension on adverse perinatal outcomes in Tigray regional state, Ethiopia: a prospective cohort study. *BMC Pregnancy Childbirth*. 2019 Dec;20(1):7.
- [19] Roberts JM, Druzin M, August PA, Gaiser RR, Bakris G, Granger JP, et al. ACOG Guidelines: Hypertension in pregnancy. American College of Obstetricians and Gynecologists. 2012.
- [20] Rana S, Lemoine E, Granger JP, Karumanchi SA. Preeclampsia: Pathophysiology, Challenges, and Perspectives. *Circ Res*. 2019 Mar;124(7):1094–112.
- [21] WHO. Drug treatment for severe hypertension in pregnancy. Geneva. 2018. 82 p.
- [22] Magee LA, Pels A, Helewa M, Rey E, von Dadelszen P; Canadian Hypertensive Disorders of Pregnancy (HDP) Working Group. Diagnosis, evaluation, and management of the hypertensive disorders of pregnancy. *Pregnancy Hypertens*. 2014 Apr;4(2):105–45.
- [23] National Guideline Alliance. Hypertension in pregnancy: diagnosis and management. *Am J Obstet Gynecol*. 2010;77(1):S1–s22.