



Letter to Editor

Covid-19: A new global threat for future generations

Corresponding Author:

Ali Akbar Oroojan;
Department of
Physiology, Faculty
of Medicine, Student
Research Committee,
Dezful University of
Medical Sciences, Dezful,
Iran.
Postal Code: 6461665145
Tel: (+98) 9166059217
Email:
aliakbar_oroojan@yahoo.com

Received 14 July 2020

Revised 22 August 2020

Accepted 9 December
2020

Production and Hosting by Knowledge E

© Oroojan *et al.* This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editor-in-Chief:

Aflatoonian Abbas M.D.

Dear Editor,

SARS-CoV-2 is a single-stranded RNA virus that acts through angiotensin-converting enzyme 2 (ACE2) as its main receptor. ACE2 expression has been revealed to be distributed in many organs of the human testis such as Leydig and Sertoli cells. Also, this virus can induce oxidative stress and increase free radicals in these cells. This event facilitates sperm DNA fragmentation that leads to reducing sperm motility and viability. As has been demonstrated, Leydig and Sertoli cells are responsible for influencing testosterone production and spermatogenesis. Angiotensin 1-7 is produced by ACE2 in the testicular tissues that activate through the Mas receptor and increase sperm motility via Phosphatidylinositol 3-kinase (PI3K)/protein kinase B (Akt) signaling pathway. In COVID-19, the expression of ACE2 in human cells is reduced. So, the Ang (1-7)/Mas/PI3K/AKT3 pathway is destroyed in the testis. This alteration in turn may decrease spermatogenesis, sperm motility, and male fertility. Moreover, this virus has been detected in the patients' semen 36 days after recovery. SARS-CoV-2 induce hyper-inflammatory syndrome by increasing inflammatory cytokines' actions through the ACE2 receptor. This event can produce pulmonary, distal alveoli, pulp, and irreversible damages in other organs that can further cause the cell apoptosis and multi-organ system failure (1-8). Therefore, due to the irreversible damage of SARS-CoV-2 to other organs such as the lungs through a mechanism similar to the mechanism of action in the testis and the long-term presence of that in the semen or male reproductive system, it can be suggested that this virus can induce male reproductive disorders in patients with COVID-19 in future. Hence, male infertility in patients with COVID-19 should be considered as a serious problem in the future generation that will require extensive clinical research to clarify its related problems and ways to treat them.

Acknowledgments

The author would like to thank the Vice Chancellor of Research, Dezful University of Medical Sciences, Dezful, Iran.

Conflict of Interest

None declared.

OPEN ACCESS

Ali Akbar Oroojan Ph.D.

Department of Physiology, Faculty of Medicine, Dezful University of Medical Sciences, Dezful, Iran.

References

- [1] Younis JS, Abassi Z, Skorecki K. Is there an impact of the COVID-19 pandemic on male fertility? The ACE2 connection. *Am J Physiol Endocrinol Metab* 2020; 318: E878–E880.
- [2] Subramanian S, Iles T, Ikramuddin S, Steer CJ. Merit of an ursodeoxycholic acid clinical trial in COVID-19 patients. *Vaccines* 2020; 8: 320–323.
- [3] Carter E, Currie ChC, Asuni A, Goldsmith R, Toon G, Horridge C, et al. The first six weeks - setting up a UK urgent dental care centre during the COVID-19 pandemic. *Br Dent J* 2020; 228: 842–848.
- [4] Frohman EM, Villemarette-Pittman NR, Melamed E, Cruz RA, Longmuir R, Varkey ThC, et al. Part I. SARS-CoV-2 triggered 'PANIC' attack in severe COVID-19. *J Neurol Sci* 2020; 415: 116936: 1–10.
- [5] Yu J, Zhang T, Zhao D, Haapasalo M, Shen Y. Characteristics of endodontic emergencies during coronavirus disease 2019 outbreak in Wuhan. *J Endod* 2020; 46: 730–735.
- [6] Anifandis G, Messini CI, Daponte A, Messinis IE. COVID-19 and fertility: A virtual reality. *Reprod Biomed Online* 2020; 41: 157–159.
- [7] Homa ShT, Vassiliou AM, Stone J, Killeen AP, Dawkins A, Xie J, et al. A comparison between two assays for measuring seminal oxidative stress and their relationship with sperm DNA fragmentation and semen parameters. *Genes* 2019; 10: 236–249.
- [8] Pal R, Bhansali A. COVID-19, diabetes mellitus and ACE2: The conundrum. *Diabetes Res Clin Pract* 2020; 162: 108132: 1–3.